

# STEWARDSHIP MANUAL

METAM SODIUM & METAM POTASSIUM



**EASTMAN**

# STEWARDSHIP MANUAL

METAM SODIUM  
& METAM POTASSIUM

**EASTMAN**



## TABLE OF CONTENTS

Table of contents .....	3
Introduction .....	5
1. Principles of soil disinfection .....	6
1.1. Soils and why they are disinfected .....	6
1.2. The basics of soil disinfection .....	7
1.2.1 Soil compartments and their role .....	7
1.2.2. Disinfectants and their transport/migration in soil .....	8
1.2.3. Factors determining disinfectant activity .....	8
1.2.3.1. Soil humidity .....	8
1.2.3.2. Soil temperature .....	8
1.2.3.3. Clay content .....	9
1.2.3.4. Organic matter content .....	9
1.2.3.5. Soil amendments .....	9
1.2.3.6. Dose rate and soil tarping .....	10
1.2.3.7. Exposure time and concentration x time concept .....	10
1.2.3.8. Recontamination of treated fields .....	11
Summary chapter 1 .....	11
2. Metam-based soil disinfectants and their major active gaseous compound .....	12
2. 1. Generalities .....	12
2.2. Characteristics of products containing metam .....	13
2.2.1. Metam sodium and metam potassium .....	13
2.2.2. Methyl isothiocyanate (MITC) .....	13
2.2.3. Biological activity as a soil disinfectant .....	13
2.3. The natural occurrence of MITC and other isothiocyanates .....	14
Summary chapter 2 .....	15

3. Mode of application of metam products in soil disinfection .....	16	5.3.4. Soil crumbling .....	33
3.1. Injection .....	16	5.4. Disinfectant application .....	33
3.1.1. Shank injection .....	16	5.4.1. Preliminary steps .....	33
3.1.2. Goose foot or blade injection .....	17	5.4.1.1. Checking the weather conditions .....	33
3.1.3. Tractor-mounted machinery illustrated .....	18	5.4.1.2. Product transfer .....	34
3.1.4. Localized point injection .....	18	5.4.2. Metam application .....	35
3.2. Drip irrigation .....	18	5.4.2.1. Soil injection .....	35
		5.4.2.2. Application by drip-irrigation .....	36
Summary chapter 3 .....	21	5.4.3. Warnings and warning signs .....	36
4. Measures that enhance metam and MITC biological activity .....	22	5.5. Re-entry period .....	37
4.1. Soil compaction .....	22	5.6. Cleaning of material and drum disposal .....	37
4.2. Water seal .....	23	5.7. Removal of plastic film seal and/or soil aeration .....	38
4.3. Plastic film tarpaulin .....	24	5.8. Cleaning of personal protection equipment (PPE) .....	39
4.3.1. Generalities and classification .....	24	5.9. General rules about the use of filter cartridges .....	39
4.3.2. Plastic film tarpaulins as part of IPM strategies .....	25	5.10. Monitoring residual MITC .....	40
4.3.3. Care of installed soil disinfection plastic films .....	25	5.10.1. Checking the work environment for safety .....	40
4.3.4. Plastic film installation modes .....	26	5.10.1.1. Photo ionisation detection (P.I.D.) .....	40
		5.10.1.2. Detection tubes .....	40
Summary chapter 4 .....	27	5.10.2. Checking for potential phytotoxicity risk after fumigation and soil aeration .....	41
5. Metam soil disinfection step-by-step stewardship .....	28	Summary chapter 5 .....	43
5.1. Storage and handling .....	28	Addenda .....	44
5.1.1. Storage .....	28	Abbreviations .....	44
5.1.2. Handling .....	28	Glossary .....	44
5.2. Check on local conditions .....	29	Legal notice .....	46
5.3. Soil preparation .....	30		
5.3.1. Removal of plant debris from foregoing crop .....	30		
5.3.2. Soil humidity .....	30		
5.3.2.1. Pre-wetting to sensitize target organisms .....	30		
5.3.2.2. Soil humidity at the time of application .....	31		
5.3.2.3. Soil humidity testing .....	31		
5.3.3. Soil temperature .....	32		

# INTRODUCTION

The storing, handling and application of soil disinfection products such as metam sodium and metam potassium requires knowledge both about the specific behaviour of these products and the volatile active gas generated following product application.

Having background knowledge of soil disinfection in general and of the factors involved enables a common-sense approach towards product use where different application techniques can be used according to specific conditions.

This manual focuses on the conditions to check and measures to take at the different stages of handling and use, including the after-treatment period. Reference is made to existing regulations and recommendations.

Each chapter contains essential information to help understand why and how general and specific measures should be observed, a detailed description of the relevant measures and a summary of highlights.

# 1. PRINCIPLES OF SOIL DISINFECTION

## 1.1. SOILS AND WHY THEY ARE DISINFECTED

Soil is a complex environment composed of mineral and organic constituents offering a biotope for beneficial as well as plant pathogenic organisms.

Mineral soil particles and organic material are present in different forms and quantities determining finer or coarser granular structure, sorption phenomena and inter- as well as intra-granular open space allowing water and gas transport.

Some soil-inhabiting organisms are beneficial, such as nitrification bacteria. However, crop pathogens (which are particularly found in cultivated soil with poor crop rotation) can cause soil-borne diseases, and soil-inhabiting crop pest organism populations such as nematodes can increase beyond crop damage threshold densities. Other types of crop-threatening organisms are weeds and their soil surviving seeds or structures. After harvest or crop removal, plant debris and plant roots that have not been totally removed enhance the risk of soil-surviving plant pathogens or pest organisms. Depending on the type of crop, different risk organisms survive at lesser or greater specific depths. A possible solution to avoid these threats is the application of soil fungicides, nematicides or herbicides.

However, only very few of the currently available plant protection products are appropriate for soil treatment. Many of them need repeated cultural treatments and present a potential danger for residue accumulation in the crop.

Soil disinfection as a pre-cultural measure is a more favourable solution. Most of the soil disinfectants have broad spectrum activity; i.e. they are often fungicidal, nematicidal and herbicidal and, depending on the application mode or technique, they can reach and treat deeper soil layers according to need.

The following table illustrates the most likely depth distribution of soil-borne pathogens and pest organisms.

## SOIL DEPTH OF VARIOUS PLANT DISEASES AND PESTS

[From: Mappes, D., 1995, *Acta Horticulturae* 382: 96-103]

SOIL DEPTH (CM)	DISEASE OR PEST ORGANISMS
0 - 20	<i>Pythium</i> spp., <i>Phytophthora citricola</i> Bacteria ( <i>Erwinia</i> , <i>Pseudomonas</i> ) Free living nematodes ( <i>Longidorus</i> , <i>Pratylenchus</i> , <i>Paratylenchus</i> )
20 - 40	<i>Sclerotium cepivorum</i> , <i>Rhizoctonia</i> spp., <i>Phoma</i> spp., <i>Didymella lycopersici</i> <i>Phytophthora fragariae</i> , <i>Verticillium albo-atrum</i> , <i>Plasmodiophora brassicae</i> <i>Thielaviopsis</i> , <i>Botrytis cinerea</i> , <i>Pyrenochaeta lycopersici</i> Root-knot nematodes ( <i>Meloidogyne</i> ) Cyst-forming nematodes ( <i>Heterodera</i> )
40 - 60	<i>Sclerotinia sclerotiorum</i> , <i>Corticium solani</i>
> 60	<i>Fusarium oxysporum</i> , <i>Rosellinia necatrix</i>

There is no sharp boundary of pest or disease occurrence at the different soil depths but the increased risk for plant disease or pest depends also on the rooting depth of the intended crop. Rooting depths of different plant species also illustrate the risk of diseased roots from a foregoing crop when left in the soil.

## ROOTING DEPTHS OF VARIOUS PLANT SPECIES

(Extended from: Mappes, D., 1995, *Acta Horticulturea* 382: 96-103)


ZONE	SOIL DEPTH (CM)	PLANT SPECIES
Shallow	0-20	Lettuce, peas, cucumbers, carrots, radishes, onions, groundnut
Medium	20-40	Potatoes, spinach, leeks, celeriac, strawberries, beans, pepper and tomatoes
Deep	> 40	Late brassicas, Brussels sprouts

## 1.2. THE BASICS OF SOIL DISINFECTION

### 1.2.1 SOIL COMPARTMENTS AND THEIR ROLE

Soil has 4 major compartments: (1) a solid mineral fraction (2) a solid organic fraction, (3) air in the inter-solids space and intra the solid soil clods, and (4) soil water forming a film around solid particles, partially filling up soil particle pores and/or freely flowing between particles.


Open channels allow easier gas diffusion and remote disinfectant activity from the point of application, although further dilution of the active gas in air will result in non-efficient concentration (see below: ct- value or concentration x time product). Increase of pore space in the soil to be disinfected is achieved by soil labour and crumbling. The charts below show the effect of soil labour or fine crumbling on pore space (light blue):



Soil labour may also give rise to a chimney effect once a disinfectant has been applied. As will be explained further on in the mode of application section, risk of premature loss of disinfectant should be reduced or even avoided by top soil layer compaction, moistening and/or soil sealing with plastic film.

Minerals such as clay and the presence of organic matter will determine sorption of the active gases. High levels of both categories of solids might necessitate increase of the effective dose rate. Soil water will dissolve disinfectants and at high presence block efficient disinfectant diffusion through the soil. This is different for the water

film surrounding the solids where the continuous exchange between the liquid and gas phase allows longer distance transport as shown below:



## 1.2.2. DISINFECTANTS AND THEIR TRANSPORT/MIGRATION IN SOIL

The distance of transport or migration of disinfectant gases in sufficient high concentration from the point of application is studied thoroughly either by chemical monitoring (gas analysers and gas sampling for further GC analysis) or, at greater depth, by introducing biological test materials (fungal cultures, nematodes, or seeds). Re-isolation of these test materials and measuring of their growth after incubation on specific media, or making counts after the needed disinfectant contact periods, will illustrate the efficacy.

Apart from the below-mentioned factors for disinfectant activity, disinfectant gas properties co-determine transport in soil as their chemical structure and molecular

size will determine sorption, solution and diffusion through micro-pores; and physical properties such as vapour pressure and gas density will determine spread in the soil atmosphere through the bigger pores and air channels.

## 1.2.3. FACTORS DETERMINING DISINFECTANT ACTIVITY

### 1.2.3.1. SOIL HUMIDITY

Humidity is needed to sensitize target organisms by making them (more) active prior to soil treatment and to regulate disinfectant diffusion processes. Sufficient humidity is needed to avoid the development of more stress-resistant survival structures, such as sclerotia or resistant spores which are less sensitive to soil disinfectants, as in the case of fungal soil borne pathogens. Best disinfectant performance is achieved within certain water holding capacity (WHC) limits. It is difficult to determine the sometimes narrow optimal value as this should be empirically determined for each plot to be treated and would change depending on the crop.

A general recommendation of soil humidity at the time of application is between 50 and 75% WHC depending on soil type and product. For metam 60% WHC is recommended.

A guideline for non-instrumental field evaluation of actual soil moisture is given further on when discussing application operational stages.

### 1.2.3.2. SOIL TEMPERATURE

Gases, and thus also disinfectants, tend to occupy the maximum available space in a closed containment environment and tend to expand more with increasing temperature. In a closed containment environment, pressure will rise with increasing temperature. The necessary volatility for a fumigating gas depends on the vapour pressure that is also temperature dependent.



Temperature also determines disinfectant solubility in soil moisture and the equilibrium of gas exchange between liquid and gas phase, as well as the adsorption and desorption phenomena at the soil solids phase.

The activity of target organisms and their sensitivity to soil disinfectants will also depend on temperature.

Adverse (too high or too low) soil temperature may lead to early morning or late afternoon/evening disinfection either to avoid risk of premature loss of the applied disinfectant at too high temperature or where it is decided to carry out delayed application to avoid low activity at too low temperature.

### 1.2.3.3. CLAY CONTENT

Clay or clay minerals have particularly strong binding capacities for most chemicals as they are often electrically charged and may show molecule-trapping spaces within their layered structure. In the case of soil disinfection, it is sometimes necessary to considerably increase dose rate (e.g. double the standard dose rate) to achieve efficient soil treatment results. Dose increase can sometimes be solved by treating only soil strips at high dose rate if the type of cropping system or field configuration allows this.

In some countries product labels specify dose rates depending on 'light' or 'heavy' soils.

### 1.2.3.4. ORGANIC MATTER CONTENT

Organic matter content in soil is mostly a result of the presence of former crops. As a standard rule, as much plant debris from former crops as possible (i.e. aerial parts and roots) should be removed before soil disinfection and next cropping. This is to avoid disinfectant loss by sorption of the gases in and on the plant debris, and

to remove remaining pest or disease inoculums in the possibly diseased foregoing crops.

It is accepted that from an organic matter content of 5%-6% onward disinfectant dose rate might need to be increased by 50%. If possible, and depending on crop type or pest/disease, strip application or treatment at a shallower depth could reduce overall dose rate to the normal rate.

Always follow local dosage regulations.

### 1.2.3.5. SOIL AMENDMENTS

Growers tend to add amendments to the soil when preparing a greenhouse or a field for cropping. The question is often raised whether it can be performed at the same time as soil disinfection.

The increase of organic matter content in the soil profile can easily be calculated; amendment at 20 ton/ha incorporated over 20 cm soil profile corresponds with approximately a 1% organic matter increase assuming soil density is approximately equal to 1.0.

Adding green manure or compost just prior to disinfection is not recommended and will necessitate increased dose rate of disinfectant when exceeding 5%-6% organic matter. Adding it for example one month before disinfectant application may allow organic material to decompose if soil temperature and moisture content are favourable. If administration of such materials is intended post-soil disinfection they should certainly not contain plant pests or pathogens.

### 1.2.3.6. DOSE RATE AND SOIL TARPING


Dose rates for soil disinfection are obtained by dose-rate studies, but for soil application they will depend moreover on the soil type and the profile (depth) of the soil to be treated with respect to target habitat and rooting depth of the intended crop.

Maximum registered dose rates may present some issues for deep treatment as they are expressed in terms of weight or volume per surface unit (kg or l/ha or g or ml/m<sup>2</sup>). In this case – as already suggested under 1.2.3.3. and 1.2.3.4. – a possible solution would be strip application.

Plastic film soil tarping or sealing can have a huge influence as the subsequent reduction of disinfectant loss during the necessary disinfection period allows reduced dose rates compared to e.g. smear rolling of the upper layer (the upper few centimeters of soil).

### 1.2.3.7. EXPOSURE TIME AND CONCENTRATION X TIME CONCEPT


Target organisms are effectively controlled when the actual disinfectant concentration is high enough during sufficient time under certain conditions of humidity and temperature. Efficacy is often expressed in terms of concentration x time product (gram per m<sup>3</sup> x hour). This could be visualized graphically by a rectangle with concentration as the height and time as the width.



Theoretical (rectangular) presentation of  $c \times t$ : green rectangles may be effective whereas blue areas are not (very high concentration during very short time or very low concentration during very long time).

In the case of MITC (methyl isothiocyanate) generation from metam, there is a quick build-up of a peak concentration of MITC followed by a decrease as MITC immediately starts to dissipate (gas expansion, sorption and degradation or metabolism). That means that presence of disinfectant either at very low concentration during a very long period or at very high concentration during a very short period may be not efficient.

The following graph illustrates the practical cases where gradual dissipation of the disinfectant is taken into account.



Concentration x time product presentation as in the case of MITC generation from metam with subsequent dissipation.

### 1.2.3.8. RECONTAMINATION OF TREATED FIELDS

Attention should be paid to avoidance of re-contamination of treated fields with pathogens such as transfer by dirt machinery or workers' boots from one field to another.

# SUMMARY CHAPTER 1

## PRINCIPLES OF SOIL DISINFECTION

- Plant pests and diseases can originate from soil. In this case they are called 'soil-borne pests and diseases'.
- The soil disinfection result or fumigant performance depends on the 'concentration x time product'; i.e. the necessity to maintain a sufficiently high concentration of the gaseous active compound during a minimum amount of time.
- Many factors affect the fumigant performance and may determine the dose-rate choice: soil humidity, soil temperature, soil clay content, soil organic matter content, organic amendments, soil pH, soil sealing.
- Basic hygiene should be respected in order to avoid recontamination of treated soils by plant pathogens.

# 2. METAM-BASED SOIL DISINFECTANTS AND THEIR MAJOR ACTIVE GASEOUS COMPOUND


## 2. 1. GENERALITIES

Metam sodium and metam potassium are both salts of N-methyldithiocarbamate. They are commercially available in aqueous solutions. Structural formulas are:

Metam sodium



Metam potassium



Applied to the soil they start decomposing into methyl isothiocyanate (MITC), carbon disulfide, hydrogen sulphide and other compounds depending on the soil pH.

MITC is the major gaseous active metam decomposition product and is known to act as a broad spectrum soil disinfectant.

Metam products are corrosive and in order to guarantee the good functioning of application devices and avoid leakage – thus guaranteeing the safe storage, handling and transfer of products – one should use the materials most appropriate for coming into contact with metam. The following table gives an overview of compatible and incompatible materials.

## OVERVIEW OF MATERIALS COMPATIBLE AND INCOMPATIBLE WITH METAM

COMPATIBLE	INCOMPATIBLE
High density polyethylene (HDPE), Polypropylene, Polyamide (Nylon 6), Polytetrafluoroethylene (PFTE; Teflon)	Copper, soft steel, aluminium, brass
Fluoro elastomere (Viton*)	Galvanized steel and zinc
Glass fibre	Polyvinylchloride (PVC)
Stainless steel	Nitrile butadiene rubber (NBR;Buna-N)
	Ethylene propylene diene monomer rubber (EPDM)
	Chlorosulfonated polyethylene rubber (CSPE; Hypalon)
	Neoprene, Butyl rubber
	Low density polyethylene (LDPE)

\*to be gradually replaced

## 2.2. CHARACTERISTICS OF PRODUCTS

### CONTAINING METAM

#### 2.2.1. METAM SODIUM AND METAM POTASSIUM

PROPERTY	METAM SODIUM	METAM POTASSIUM
Content of active ingredient	510 g/L or 42.1% w/w	690 g/L or 54% w/w
Formulation type (code)	SL (miscible with water)	SL (miscible with water)
Vapour pressure of the active ingredient	$5.75 \times 10^{-2}$ Pa at 25°C Moderately volatile	Moderately volatile
Volatility from water (Henry's law constant)	$8.34 \times 10^{-6}$ Pa.m <sup>3</sup> /mol at 20°C Very slightly volatile from water	
Decomposition temperature of the active ingredient	150°C	150°C
Shelf life of formulation	2 years at ambient temperature	2 years at ambient temperature
Dilution stability	Stable after accelerated storage (14 days at 54°C)	

#### 2.2.2. METHYL ISOTHIOCYANATE (MITC)

PROPERTY	VALUE
Vapour pressure	1739 Pa at 20°C Highly volatile substance
Solubility in water	8.94 g/L at 20°C and pH 7.5 Readily soluble in water
Volatility from water (Henry's law constant)	14.2 Pa.m <sup>3</sup> /mol at 20°C Moderately volatile from water
Gas density (air=1.0)	2.5

The overviews of properties demonstrate the favourable stability, solubility and volatility behaviour of the metam products as well as the disinfectant capability of the generated active MITC.

#### 2.2.3. BIOLOGICAL ACTIVITY AS A SOIL DISINFECTANT

Metam products are broad spectrum soil disinfectants. The registered activities cover the large group of nematodes, fungi and weeds.

Metam and MITC are also known to control certain stages of soil-borne pest insects.

Examples of different species among these groups are summarized in the following table.

NEMATODES	FUNGI	WEEDS
<b>Root-knot nematodes:</b>	<i>Fusarium</i> spp.	<i>Amaranthus</i> spp.
- <i>Meloidogyne</i> spp.	<i>Phialophora</i> spp.	<i>Galium aparine</i>
<b>Cyst nematodes:</b>	<i>Phoma</i> spp.	<i>Malva</i> spp.
- <i>Globodera</i> spp.	<i>Phytophthora</i> spp.	<i>Matricaria</i> spp.
- <i>Heterodera</i> spp.	<i>Pythium</i> spp.	<i>Mercurialis annua</i>
<b>Free living nematodes:</b>	<i>Rhizoctonia</i> spp.	<i>Poa annua</i>
- <i>Paratylenchus</i> spp.	<i>Sclerotinia</i> spp.	<i>Senecio</i> spp.
- <i>Pratylenchus</i> spp.	<i>Verticillium</i> spp.	<i>Solanum</i> spp.
- <i>Rotylenchus</i> spp.	<i>Aphanomyces</i> spp.	<i>Sorghum halepense</i>
- <i>Trichodorus</i> spp.	<i>Macrophomina</i> spp.	<i>Stellaria</i> spp.
<b>Stem nematodes:</b>	<i>Sclerotium</i> spp.	<i>Taraxacum officinale</i>
- <i>Ditylenchus</i> spp.	<i>Monosporascus</i> spp.	<i>Portulaca</i> spp.
- <i>Aphelencooides</i> spp.	<i>Thielaviopsis</i> spp.	<i>Orobanche</i> spp.
		<i>Cuscuta</i> spp.
		<i>Echinochloa</i> spp.
		<i>Polygonum</i> spp.
		<i>Chenopodium</i> spp.
		<i>Cirsium</i> spp.

## 2.3. THE NATURAL OCCURRENCE OF

### MITC AND OTHER ISOTHIOCYANATES

MITC and many other isothiocyanates (ITCs) are generated by maceration of specific crops (e.g. *Brassicaceae*) conducting enzymatic (myrosinase) conversion of glucosinolates into corresponding ITCs. Many of them show biological activity. This is also the basis of biodisinfection, i.e. growing appropriate crops on the field to be treated, followed by incorporation of the fully grown crop. Alternatively, the crop can be harvested and spread and incorporated at another field location. The problem is that glycosinolate content may fluctuate and thus the content of the active compound is not well known.

MITC is known to be generated from its precursor glucoapparin particularly in capers (*Capparis spinosa*), horseradish (*Armoraria rusticana*), *Cleome spinosa* (Spider flower) and in the seeds of other species of the genus *Cleome*.

A few references from literature are:

Kjaer A., 1960. Naturally derived isothiocyanates (mustard oils) and their parent glucosides. Fortschr. Chem. Org. NatStoffe 18: 122-176.

Ahmed Z.F., Rizk A.M., Hammouda F.M. and Seif El-Nasr M.M., 1972. Glucosinolates of Egyptian *Capparis* species. Phytochemistry 11: 251-256.

Matthäus B. and Özcan M., 2002. Glucosinolate composition of young shoots and flower buds of capers (*Capparis* species) growing wild in Turkey. J. Agric. Food Chem. 50 (25): 7323-7325.

Kaur R., Rampal G. and Pal Vig A., 2011. Evaluation of antifungal and antioxidative potential of hydrolytic products of glucosinolates from some members of Brassicaceae family. Journal of Plant Breeding and Crop Science 3(10): 218-228.

The latter publication of 2011 reports on up to 12 different glucosinolates (precursors of ITCs) with content between 6.55  $\mu\text{mol/g}$  in raw flower buds of *Capparis spinosa* (i.e. the edible capers) and up to 45.56  $\mu\text{mol/g}$  in young shoots of *Capparis ovata*. About 90% of the total glucosinolates found is glucoapparin, the precursor of MITC.

It is also interesting to know that human consumption of many types of edible crops, leads to the generation of ITCs from the plant glucosinolates and the absorption of ITCs in the digestive tract. Moreover, diets including the associated crops such as Brussels sprouts, brassicas, cress etc., are known to exert anti-cancer activity in mammals – a phenomenon which is being increasingly recognized.



## SUMMARY CHAPTER 2

### METAM-BASED SOIL DISINFECTANTS AND THEIR MAJOR ACTIVE GASEOUS COMPOUND

- Metam products are available either as aqueous solutions of sodium or potassium salts (code SL (Soluble Liquid)).
- Both formulations are stable at room temperature and once applied in soil they generate the more volatile methyl isothiocyanate (MITC) as the most biologically active decomposition compound.
- MITC shows solubility and volatility characteristics suitable to act as a soil fumigant.
- MITC and other isothiocyanates are known either to occur naturally or to be enzymatically generated in damaged or macerated plant species.

# 3. MODE OF APPLICATION OF METAM PRODUCTS IN SOIL DISINFECTION

The goal of this chapter is to make the most appropriate choice of application mode according to local conditions and available equipment.

The choice for a metam application technique can depend on different factors:

- Area or surface to be treated
- Open field or protected area (greenhouse, tunnel)
- Presence of drip irrigation installation
- Distance from residential zones
- Local uses of professional applicators

The types of techniques are:

- Soil injection
- Drip irrigation

## 3.1. INJECTION

The principle is the deep application of the soil disinfectant mainly either by shank injection or goose foot injection.

The advantage of deeper soil profile treatment is that it allows a faster and more homogenous application.

### 3.1.1. SHANK INJECTION

Shanks are knife shaped and the applied liquid disinfectant escapes through the tubing opening at the deep end of the shank. The configuration preferably contains an anti-drip border in order to avoid dripping when the application machine is lifted out of the soil, e.g. when moving to another strip to be treated.

The shank position on the machinery can be adjusted according to soil type and target application depth. This means that for heavy soils the shanks' interspacing could be smaller. For deep injection or more homogenous soil profile spread, the depth positioning of shanks could be adjusted or shank depth could alternate.

The following pictures illustrate a variety of shanks.







Shallow shanks for greenhouse use



Various types of goose-foot blade injection system

### 3.1.2. GOOSE FOOT OR BLADE INJECTION


This configuration is very suitable for heavy soils and big open field areas. As the tractor-mounted machine advances, the goose-foot shaped blade lifts the soil while the disinfectant piping exit underneath allows good spread of the liquid disinfectant on the soil at the chosen depth under the entire surface of the blade if equipped with a spraying nozzle.

The following pictures illustrate this type of application machine as well as different types of goose-foot blades. The presence of a spading section after the injection part (as shown top right) homogenizes the treated soil.



Various types of goose-foot blade injection system

Highly successful commercially produced machine versions are shown below:



- **Mix Tiller Deeper soil disinfection machine** (manufactured by Forigo)

- **Rotary spading machine with injection** (manufactured by Imants)



### 3.1.3. TRACTOR-MOUNTED MACHINERY ILLUSTRATED



Back-mounted tank



Front-mounted tank



High-volume tanks



### 3.1.4. LOCALIZED POINT INJECTION


A very specific application device is a manual point deep injector. This application method is typically used in vineyards for the control of *Armillaria mellea* fungus, the causal agent for mushroom root rot, and *Xiphinema* spp. nematodes that are vectors of the grapevine fan leaf virus (GFVL) disease.




### 3.2. DRIP IRRIGATION

A very common method is drip irrigation, which is employed in many high-value crop greenhouses and fields. Drip irrigation tubes are spaced 20 to 50 cm apart and perforated every 15 to 30 cm. They should be buried at 5 cm soil depth or covered by plastic film soil tarpaulin. Metam is then applied as a water-diluted solution/mixture where the metam is supplied by an automatic dosage controller/dosing pump such as e.g. the Dosatron. A dilution of 0.1 to 2.0% should be achieved. The system must be fitted with an anti-return valve. It takes 1 to 4 hours to apply 10 to 40 mm of dilution. In order to obtain good metam solution diffusion, the soil should be reasonably compacted (especially in the case of sandy soil).

#### Metam Diffusion



## EXAMPLE OF MOBILE INJECTION SYSTEM



## EXAMPLE OF FIXED DRIP IRRIGATION INSTALLATION

(Photos courtesy Biotek Ag Spain)



Irrigation house



Central control unit to program irrigation time and flow



Metam is introduced at the head of the irrigation system by pouring the product into the tank made available for this purpose (not such a common method)



Alternatively, a pump can be connected directly to the drum containing metam (most common method)



A) Pump powered by electricity

B) Pump powered by the water flow



When there is no injection pump available, the product can be incorporated into the water by suction through a Venturi system



The system can be regulated to maintain the desired concentration of metam in water

## SUMMARY CHAPTER 3

### MODE OF APPLICATION OF METAM PRODUCTS IN SOIL DISINFECTION

- Metam can be applied by two methods: soil injection and drip irrigation system.
  - Injection allows the product to be applied at 10 to 40 cm deep with machines equipped with shanks or goose foot-shaped blades. The machines are mounted at the rear of a tractor. Localized injection for small areas can be done with a manual point deep injector.
  - Application through the existing drip irrigation system is done by incorporating Metam in the irrigation water with the help of a dosing pump.

# 4. MEASURES THAT ENHANCE METAM AND MITC BIOLOGICAL ACTIVITY

This chapter describes the measures that need to be taken to optimize successful soil disinfection.

From the 'concentration x time' product concept (see chapter 1.2.3.7) a longer disinfectant contact time combined with disinfectant concentration will give a better result.

The principle is to avoid premature loss of the applied chemical. This can be achieved by quick homogenization of the applied metam in the soil and sealing of the soil. Another important factor is the observation of recommended temperature limits for application: at too low a temperature the product will not work (too low a volatility of the active compound and inactivity or thus lower sensitivity of the target organisms) and at too high a temperature the active compound will dissipate or be lost by too fast an emission. High emission rates should also be avoided in the interests of worker, bystander and resident safety.

For soil sealing three major techniques or combinations thereof are available:

- Soil compaction
- Water seal
- Plastic film tarpaulin

## 4.1. SOIL COMPACTION

Soil compaction is often inherent to the application by mechanical soil injection through the presence of a compaction roll on the application machine.



A view of the soil surface after passage of a compaction roll

The compaction roll is mounted directly behind the rotavating machinery or the spading section and rotates in opposite motion to the machine in order to get a smooth, sometimes mirror-like, surface on fine crumbled soil. The upper 1-2 centimetres of soil is very compacted and slows down MITC emission. On more sophisticated models the roller is driven hydraulically, and pressure and rotation speed can be programmed. An alternative to flat compaction is the compactor/bed-shaper. In some applications the additional wetting of the top soil increases barrier properties. Additional wetting should be avoided when soil is too clayish because completely blocked soil pores may reduce product efficacy in the top layer.

The following pictures illustrate some commercial versions of the application machine fitted with compacting rollers.



Soil compacting rolls on IMANTS machines (left) and on a FORIGO machine (right)

## 4.2. WATER SEAL


Wetting of the upper few centimetres of soil will reduce premature disinfectant loss. As indicated under 4.1., too-clayish soil is less appropriate as completely blocked soil pores may reduce product efficacy in the top layer.

Under warm or hot conditions, or wind in open field, a dry crust may quickly form, giving rise to cracks and loss of its disinfectant-emission-reducing qualities. Intermittent irrigation could offer a solution, as illustrated by the following results.

Shank injection studies were performed to study off-gassing rates of MITC following metam application. The following table summarizes intermittent irrigation times and quantities at the Lost Hills site (Merricks, L.D., 2001, Agrisearch study) and single irrigation time and quantity at the Kern site (Merricks, L.D., 2002, Agrisearch study).

IRRIGATION SESSION		1	2	3	4	5
TIME BETWEEN METAM APPLICATION & IRRIGATION SESSION		4 hours	12 hours	16 hours	24+4 hours	24+12 hours
INTERMITTANT IRRIGATION	QUANTITY OF WATER	10 mm	5 mm	5 mm	5 mm	5 mm
	TIME OF IRRIGATION	11:00 – 13:00	19:00 – 20:00	23:00 – 24:00	11:00 – 12:00	19:00 – 20:00
SINGLE IRRIGATION	QUANTITY OF WATER	20 mm				
	TIME OF IRRIGATION	11:00 – 13:00				

Results showing low off-gassing of MITC from metam-treated soil by intermittent irrigation at the Lost Hills site as compared to single irrigation at the Kern site are summarized in the following graph:



**Comparison of MITC flux to the atmosphere during 96-hour monitoring period with 24 samplings with 4 hour-intervals between intermittent irrigated and single irrigated metam-treated fields**

## 4.3. PLASTIC FILM TARPAULIN

### 4.3.1. GENERALITIES AND CLASSIFICATION

Plastic film can be applied to improve efficacy while simultaneously reducing disinfectant emission risks. Standard low density polyethylene (LDPE) film of 20-30 µm thickness is widely used. Although LDPE presents a certain permeability to gases, the use of such films helps greatly to reduce the MITC emissions, in two main ways. Firstly, the plastic tarpaulin will prevent the upper layer of the soil from drying and thereby reduce upward water and MITC movement that would result in more MITC escaping from the soil surface. Secondly, MITC being highly soluble in water, the fraction that escapes from the soil surface will be trapped by the water present in the form of film and droplets on the inner plastic tarpaulin surface. MITC may ultimately return to the soil.

When used in open field, another very good reason to use plastic film is that it can combine the soil disinfection action of metam with that of the solarization (see section 4.3.2.)

There is a large variety of plastic film for soil disinfection available on the market globally. While criteria such as mechanical resistance and workability are imperative, the gas barrier properties are the most important for an optimal treatment result. Unfortunately, the more gastight plastic films are more expensive.

The so called V.I.F., T.I.F. and F.I.F. films often have a 3-, 5- or even 7-layered structure where the central layer, often of only a few micrometers' thickness, is the real gas barrier layer, surrounded by adhesive layers and layers present simply for mechanical resistance of the embedded barrier. The total film thickness is often in the 35-40 µm range. An overview of all commercially available film is out of the scope of this manual but here follows a useful classification table.



CLASS	ABBREVIATION	BARRIER COMPOUNDS
Not completely impermeable		LDPE, PVC, biodegradable starch and polylactic acid (PLA) based films, metal oxide coatings
Virtually impermeable films	V.I.F.	ethylene vinyl alcohol (EVOH), polyamide (PA) and others
Totally impermeable films	T.I.F.	
Fully impermeable films	F.I.F.	

Biodegradable film needs to be about double the thickness of LDPE for comparable barrier properties.

Although plastic film sealing for soil disinfection is very useful from a safety and dose-rate reduction perspective, it represents an extra application cost as well as removal and waste treatment costs. Cleaning or washing of used soil disinfection film is not easy. However recycling is possible and worthwhile with the higher grade impermeable film containing the more expensive gas barrier compounds.

### 4.3.2. PLASTIC FILM TARPAULINS AS PART OF IPM STRATEGIES

In more southern regions (such as the South of France, Mediterranean countries and their isles, and Northern Africa), soil disinfection can be combined with soil

solarization as part of integrated pest management (IPM) strategies where the disinfectant action considerably reduces or weakens target organism populations for consecutive control by the longer solarization phase. Analogously, pre-cultural soil disinfection can precede a later application of soil-applied antagonistic preparations.

Plastic films for soil solarization should contain ultraviolet blockers in order to offer longer resistance to decay.

### 4.3.3. CARE OF INSTALLED SOIL DISINFECTION PLASTIC FILMS

Applicators/workers and farmers/growers should be aware that the smallest pinhole, especially in the more expensive impermeable film types, considerably reduces the active concentration of disinfectant over the underlying soil surface.

It is imperative to remove from the soil, as far as possible, all sharp stones and plant materials that could harm the plastic film.

Stepping on an installed film should be avoided as it increases the risk of pinholes, especially if stones or other sharp materials are left in the soil. A footstep upon the film on top of the relatively loose soil will stretch out the film and considerably reduce the local thickness, giving rise to a less gas impermeable zone.

It is useful to apply some water on top of the installed film, not only to reduce the space between soil and film to a minimum but also to avoid tearing of the film in outdoor treated fields due to the wind. In greenhouses, sprinklers could be used for the first purpose.

Sometimes soil is thrown on top of the film in different places but this raises the risk of pinholes.

#### 4.3.4. PLASTIC FILM INSTALLATION MODES

There are 2 possibilities:

- The film is installed in one run with the combined application and homogenizing machine as shown in the following pictures:



Single pass equipment:  
application/incorporation/plastic film sealing machine

- In the case of drip-irrigation product application, plastic film is applied over the drip lines prior to the disinfectant administration



Indoor drip irrigation



Outdoor drip irrigation

If manually unrolling and digging the film is necessary, workers should take all necessary precautions to protect themselves from exposure including the use of Personal Protective Equipment (PPE) and should not work soon after the passage of the application machine. Additional information regarding PPE and re-entry are provided in subsequent sections of this manual.

# SUMMARY

## CHAPTER 4

### MEASURES THAT ENHANCE METAM AND MITC BIOLOGICAL ACTIVITY

- The efficacy of applied metam as a soil disinfectant can be increased by avoidance of premature loss of the generated MITC.
- Soil sealing measures can be summarized under 3 types:
  - soil compaction
  - water sealing (preferably with inter-mittent irrigation)
  - plastic film sealing (with good gas barrier properties)
- It is compulsory to carry out one or combined measures immediately after disinfectant application, also from a worker-, bystander- and resident-safety perspective.
- The application of plastic film enables additional solarization effects under appropriate climatic conditions.
- In order to optimize plastic film sealing performance extreme care should be taken not to damage film by sharp objects or footsteps.



# 5. METAM SOIL DISINFECTION STEP-BY- STEP STEWARDSHIP

This chapter aims to provide the necessary information for optimizing the soil disinfection result and to preserve safety at work and minimize impact on the environment. It includes information on storage and handling, checking local conditions, soil preparation, the different application purposes and particular situations, and after-care and monitoring. **Always keep the Material Safety Data Sheet close at hand.**

## 5.1. STORAGE AND HANDLING

### 5.1.1. STORAGE

Metam products should always be stored in closed original packaging showing original label and applicable hazard logos. The storage room should be cool and ventilated and should be kept locked at all times. It should be located far from buildings with human activity and habitations, including those for animals. The maximum volumes to be stored at one site should correspond with local regulations.

Apart from the original plastic drums, nothing other than stainless steel metal containers should be used for storage. Empty packaging should not be reused.

The storage area should be inaccessible to children and anybody else not involved in the farm operations, and display no-entry signs and signs indicating the presence of hazardous/corrosive materials.

Regarding the construction of the room, it should have a basin-shaped floor in case of product leakage, and roof and wall materials should prevent the stored products from heating up. The temperature should not exceed 35°C.

### 5.1.2. HANDLING

Product handling is the most critical phase.

The choice of material is important for the product transfer connecting systems. A list of metam-compatible and incompatible materials was given under 2.1.

Before starting any handling the worker should dress in the appropriate personal protection equipment (PPE). This covers both clothing and protection apparatuses:

- Eye protection goggles
- Rubber gloves
- Product resistant synthetic coverall
- Rubber boots
- Respiratory mask with a canister approved for organic vapour with a boiling point higher than 65°C. Combined filters such as A2B2-P3 are filled with activated carbon impregnated with a gas molecule retaining substance. The filter protects for example against organic and inorganic gases and vapour (max 5000 ppm) as well as against toxic substances. It is for general use against plant protection chemicals. In a confined space, a self contained breathing apparatus (SCBA) with full face mask is recommended

Air purifying respirators (APRs) are available in both half-face and full- face models.

●  
Half-face  
APR



●  
Full-face  
APR

When filling the application machine tank, leak-free connections should be used. Avoid any spill, drain or drip to soil.

On the transfer site, the floor should be impermeable in order to be able to collect product in case of spillage. Risk of evacuation to drains and ditches is prohibited. Never perform such handling close to waterways or ditches.

## 5.2. CHECK ON LOCAL CONDITIONS

Prior to application the contractor needs to know or check the local conditions of the greenhouse or field to be disinfected.

A checklist should be used to assess the necessary measures to take, and help to decide on execution in the case of temporary adverse conditions.

The following overview could serve as a checklist:

POINT NUMBER	DESCRIPTION	COMMENTS	TICK BOX
1	Localization	Check for neighbouring buildings (public and habitations), presence of water surfaces, draining ditches, animal husbandry, adjacent crops	✓
2	Field condition	Check for presence of plant debris, appropriate soil preparation (crumbled and pre-wetting for target organisms sensitization) soil temperature and soil humidity	✓
3	Greenhouse	Check for broken windows, cracked or torn plastic roof or walls	✓
4	Wind	Check for wind direction and consult forecast including for the days immediately following	✓
5	Temperature	Consult temperature forecast and the risk of temperature inversion* including for the days immediately following	✓

\*Temperature inversion is a meteorological situation where temperature increases in higher air layers with lower temperature in the lower layer creating a trap for pollutants by preventing their dilution into the atmosphere.

Prior to disinfection the weather forecast for the day of application and the 48-hour period following the disinfection must be checked to determine if unfavourable weather conditions exist or are predicted and whether disinfection should proceed (refer to section 5.4.1.1.).

Of course, the grower and the operator may each have their own responsibility, especially in carrying out correct soil conditioning favourable to soil disinfection.

If necessary the application should be postponed until conditions are more favourable.

To protect aquatic organisms:

- do not apply to drained soils.
- in fields vulnerable to surface runoff, respect an untreated vegetative buffer zone of 10 m to surface water bodies.

## 5.3. SOIL PREPARATION

### 5.3.1. REMOVAL OF PLANT DEBRIS FROM FOREGOING CROP

As explained in the basics on soil disinfection (1.2.3.4.), the target field to be treated should be as free as possible of plant debris as it could raise the organic matter content to a critical level or present a source for re-infection of the soil. Particular attention should be given to the removal of deep-rooting crops.

### 5.3.2. SOIL HUMIDITY

As indicated under 1.2.3.1. there are two main reasons to check and if necessary adjust soil humidity.

### 5.3.2.1. PRE-WETTING TO SENSITIZE TARGET ORGANISMS

Soil moisture content should be high enough to activate the target organisms responsible for soil-borne plant pests and diseases and weed seeds. If soil humidity adjustment is necessary, the period prior to soil disinfection may be 5 to 10 days' long depending on the type of target organisms and the resistance of inactive surviving structures or stages, as well as the ambient temperature influencing the speed of reactivation after soil wetting. In outdoor situations, the soil to be disinfected may have a sufficiently high moisture content level depending on recent rainfall.

Where possible or available, sprinkler or drip irrigation is the easiest way to prepare the soil for this purpose. Outdoors and for large surface areas, mobile sprinklers can be used. In some cases, when soil humidity is as high as 15 cm below surface, soil tillage could mix up with the dryer top layer. The latter technique could also be used if soil moisture needs to be adjusted again at the time of application (5.3.2.2.)

### 5.3.2.2. SOIL HUMIDITY AT THE TIME OF APPLICATION

In contrast to soil pre-wetting for the sensitization of target organisms, soil moisture content for the product application itself is more critical from a gas diffusion perspective and for sorption processes during soil disinfection. It is compulsory to perform soil humidity testing just before application in order to adjust if necessary.

### 5.3.2.3. SOIL HUMIDITY TESTING

Field testing is preferably done by measurement and can be confirmed by a simple 'feel and appearance' method.

A well-known quick method for not-too-extreme soil types consists of taking a firm lump of soil that should easily crumble into pieces when it is then dropped back onto the ground.

The most optimal humidity level lies between 50 and 75% (even beyond 75% in case of bed-shaping) of the water-holding capacity. The most accurate measurement method is by laboratory weighing and drying but this is time consuming. A more practical option is an electronic device that can be used in the field.

Instruments for soil humidity reading are often based on electrical resistance or on soil suction (pF or water potential) measurement. Read-out is then respectively absolute moisture content in % or in pF/centibar/MPa. For a soil with known % water retention capacity, % relative moisture content can be calculated. Other instruments contain an arbitrary scale that could be compared and roughly calibrated for %.

A detailed overview of how to use 'feel and appearance' criteria to estimate soil moisture can be found in the following overview from the USDA National Resource Conservation Service:

The feel and appearance of soil vary with texture and moisture content. Soil moisture conditions can be estimated, with experience, to an accuracy of about 5 percent. Soil moisture is typically sampled in 30 cm increments to the root depth of the crop at three or more sites per field. It is best to vary the number of sample sites and depths according to crop, field size, soil texture, and soil stratification. For each sample the "feel and appearance method" involves:

- Obtaining a soil sample at the selected depth using a probe, auger, or shovel
- Squeezing the soil sample firmly in your hand several times to form an irregularly shaped "ball"
- Squeezing the soil sample out of your hand between thumb and forefinger to form a ribbon

- Observing soil texture, ability to ribbon, firmness and surface roughness of the ball, water glistening, loose soil particles, soil/water staining on fingers, and soil colour. [Note: A very weak ball will disintegrate with one tap of the hand. A weak ball disintegrates with two to three taps]
- Comparing observations with photographs and/or charts to estimate percent water available and the water level depleted below field capacity

#### Appearance of different types of soil at various moisture conditions:

- Fine sand and loamy fine sand soils
- Sandy loam and fine sandy loam soils
- Sandy clay loam and loam soils
- Clay, clay loam, and silty clay loam soils

Details of each above-listed soil category with accompanying pictures are included in a document titled Estimating Soil Moisture by Feel and Appearance which can be found at the following link (active at time of publication):

<http://msue.anr.msu.edu/uploads/235/67987/lyndon/FeelSoil.pdf>

The following table also gives a useful overview of different soil types:

### SOIL MOISTURE IDENTIFICATION CHART

[R.W. Harris and R.H. Coppock (Eds.), 1978. University of California Division of Ag Science leaflet 2976]

% MOISTURE	SAND	SANDY LOAM	CLAY LOAM	CLAY
Close to 0%	Dry, loose, single grained flows through fingers	Dry, loose flows through fingers	Dry clods breakdown into powdery condition	Hard, baked, cracked surface, loose crumbs on surface
50% or less	Appears dry, will not form ball	Appears dry, will not form ball	Crumbly, holds together with pressure	Pliable, will ball under pressure
50%–70%	Same as above	Will ball, but will not hold together	Forms a ball, slight slick with pressure	Forms a ball ribbon between fingers
75% to field capacity	Sticks together, forms a weak ball	Forms a weak ball, will not become slick	Forms a ball very pliable, readily forms a slick	Easily ribbons between fingers
Field capacity	Under pressure, moisture appears on hand	Same as sand	Same as sand	Same as sand

### 5.3.3. SOIL TEMPERATURE

Soil temperature prior to soil disinfection, together with soil humidity, affects the target organisms' sensitivity. As a basic rule for metam soil disinfection, a minimum



temperature of 10°C and a maximum temperature of 25°C are required. The average temperature during the disinfection period determines the minimum number of days before proceeding to soil aeration prior to sowing or planting.

Soil temperature should be checked at 10-15 cm depth. Both traditional and electronically operated thermometers are commercially available and can be used for this purpose.

Soil temperature is not always under control, especially in outdoor conditions.

For both outdoor and indoor disinfection, alternate timing of the application in the early morning or in the evening could meet the temperature requirements.

For soil disinfection under protection, soil and air temperature can sometimes be controlled. This would apply both if the pre-wetted period and disinfection conditions were too cold or too high for optimal activity.

However, in cases where soil and air temperature are too low, expensive energy is needed to reach or to exceed lowest temperature requirements.

In the case of too high temperature, risk application timing can be considered and in case of availability, the use of shadow screens could help to solve such issues.

#### 5.3.4. SOIL CRUMBLING

Homogenization of the soil layer at least over the soil profile to be disinfected can be performed during the plant debris removal operation or when re-wetting or adjusting soil moisture content. The double goal is to obtain suitable homogenous sensitization and to avoid the presence of gas impenetrable soil clods that would limit the control of enveloped target organisms.

Large clods can reduce soil sealing by causing a chimney effect.

The mechanical tools used for soil crumbling are spading and rotavating machinery preferably used several days prior to the disinfectant application for the above-mentioned target sensitization reasons.

It is pointless to carry out soil labour on non-disturbed soil shortly prior to soil disinfection.

## 5.4. DISINFECTANT APPLICATION

Safety measures to be taken and the required personal protection equipment will depend on the type of handling and application activity and are discussed in more detail in the Safety Data Sheet (SDS) and under the respective sections below. **Always keep Safety Data Sheets available in case a spill or other incident occurs.**

### 5.4.1. PRELIMINARY STEPS

As emphasized under 5.2. the checking of local conditions is compulsory and may be the deciding factor with regard to method of application. However, the check of the product transfer, application equipment and safety equipment shortly before use is just as important.

Good Agricultural Practices (GAP) must be followed during all disinfectant applications.

#### 5.4.1.1. CHECKING THE WEATHER CONDITIONS

Do not apply in adverse weather conditions.

**Conditions to avoid:**

- The presence of temperature inversion or air-stagnation in the area where the disinfection is planned
- Strong winds, heavy rain or thunderstorms in the application phase and in the hours immediately following (check the weather forecast)
- The phenomena of drift gas where the air mass moves in unpredictable directions even at considerable distances from the treated site

When disinfecting field plots upstream of waterways, ponds, water wells, etc. retention works should be foreseen downhill by making soil heaps or a basin.

**5.4.1.2. PRODUCT TRANSFER**

Metam transfer from drum to tank should be carried out with appropriate connections remote from water surfaces, ditches and habitations. The handlers' personal protection equipment (PPE) should fulfil general and local requirements.

All tanks, hoses, fittings, valves and connections must be serviceable, tightened, sealed and not leaking.

Dry disconnect couplings (closed transfer system) must be installed on all tanks and transfer hoses.

Sight and pressure gauges must function properly.

Tanks, hoses and fittings should be designed to withstand the pressure of the system and be resistant to metam (For the compatible and not-compatible materials see section 2.1.).

Packaging pressurization for emptying operations:

- **IBCs (Intermediate Bulk Containers of about 1000 L):** we recommend emptying the container by gravity through the bottom outlet valve, without pressure. If pressure is applied inside the container, the overpressure should not exceed 40 mbar (0.04 bar)
- **Drums:** overpressure inside the drums should not exceed 150 mbar (0.15 bars)

Replace drum closure lids after transfer, even on empty drums. For disposal/recycling of empty drums see section 5.6.

**PERSONAL PROTECTION EQUIPMENT (PPE)  
BY PRODUCT TRANSFER**

- Chemical-resistant rubber gloves
- Chemical-resistant rubber boots
- Synthetic coverall (trousers preferably slipped over the rubber boots to avoid entry of liquids)
- Eye or facial protection
- Air Purifying Respirator (APR): A2B2-P3 or similar filter cartridges mounted on half-face mask or preferably full-face mask
- In a closed environment self-contained breathing apparatus (SCBA) is preferred

## 5.4.2. METAM APPLICATION

Here a distinction should be made between the two major application methods:

1. Soil injection
2. Drip irrigation

Depending on the tractor type used, indoor or outdoor application, drip irrigation and the activities of the workers involved, PPE may vary.

### 5.4.2.1. SOIL INJECTION

Only tractors with closed cabins may be used for the mechanical incorporation of metam.


The machine can either be equipped with a smooth roll at the rear in order to form an even and smooth upper soil layer or with a plastic film application device. The aim of both systems is to reduce the gas emission in the air, to reduce bystander and operator exposure and at the same time to increase the soil disinfection efficacy.

It is not advisable to use a tractor with open or no cabin for the application of metam products.

#### Machine check prior to application:

- Application equipment must be in good working order
- Sight gauges and pressure gauges must function properly
- Nozzles or shanks and metering devices must be of correct size, sealed and unobstructed
- Each nozzle/shank must be equipped with a flow monitor (this could be mechanical, electronic or 'red-ball' type)

The following pictures illustrate a few different types of flow monitoring equipment.



Multi-channel flow meter



Single flow meter



Electronic flow controller for individual product release channels (TeeJet)

Before using a disinfection rig for the first time, or when preparing it for use after storage, the operator must check the following items carefully:

- Check and clean or replace the filter element if necessary
- Check all tubes and chisels/shanks to make sure they are free of debris and obstructions
- Check and clean the orifice plates

Injectors must be below the soil surface before product flow begins. Each injection line must either have a check valve located as close as possible to the final injection point, or drain/purge the line of any remaining disinfectant prior to lifting injection shanks from the ground.

Do not lift injection shanks from the soil until the shut-off valve has been closed and the disinfectant has been depressurized. Remaining product can be passively drained or actively purged by compressed air.

### PERSONAL PROTECTION EQUIPMENT (PPE) IN CASE OF MECHANICAL INCORPORATION

Only tractors with closed cabins may be used:

It is advisable to use closed cabins of Category 4 (EU Standard EN 15695-1). It will protect the operator in the cabin from vapours, aerosols and dust.

- Rubber boots
- Cotton coverall

In case of other type of cabins (Categories 1, 2 or 3), the following PPE must be worn in addition to those listed above:

- Facial mask with active carbon filter type A (brown colour code) for gases and organic vapours with boiling point > 65°C

It is important not to bring any clothes, boots or devices spoiled with metam into the cabin and to wear the above-mentioned PPE equipment for any intervention on the field or the application machine.

When applying the product indoors it is strongly recommended to keep greenhouses or tunnels closed until venting.

The most critical phases are the coupling and uncoupling of the metam drum or container to the drip-irrigation device/pump. It is therefore recommended that operators be equipped the same way as for product transfer from drum or container to an application machine tank, as shown in the box below.:

### PERSONAL PROTECTION EQUIPMENT (PPE) IN CASE OF DRIP DISINFECTION

- Chemical resistant rubber gloves
- Chemical resistant rubber boots
- Synthetic coverall (trousers preferably slipped over the rubber boots to avoid liquids entry)
- Eye or facial protection
- Air Purifying Respirator (APR): A2B2-P3 or similar filter cartridges mounted on half face mask or preferably full face mask

#### 5.4.2.2. APPLICATION BY DRIP-IRRIGATION

The application itself does not necessitate close exposure of the operator once the system is coupled to the metam source.

Prior to product handling the water supply metering system, dilution pump, no-return valve free and drip lines should all be checked for correct functioning. The main water pipe should be free of obstruction.

#### 5.4.3. WARNINGS AND WARNING SIGNS

We strongly advise growers that prior to every campaign they should inform residents, within a radius of 200 meters of the plots to be treated, about the periods of application. Treated fields and greenhouses should be indicated by no-entry warning signs.

Greenhouse entrance doors are advised to be kept closed until ventilation. 'No trespassing' or 'Disinfection in progress' signage could be used.



## 5.5. RE-ENTRY PERIOD

Due to the volatile chemical properties of metam and MITC and the potential for worker exposure, entry into a treated field is restricted during a certain period of time to handlers wearing appropriate PPE. Re-entry period is a specific amount of time during which entry into treated fields by anyone other than a trained handler, dressed in the correct PPE, is prohibited.

**The re-entry period starts from the end of the product application.** The length of the re-entry period depends on whether the application was done in open field or in greenhouse.

### Open field

Depending on the climate of the region, the recommendation is to wait minimum 7 days before entering a treated field. Always follow local label requirements.

### Greenhouse

The re-entry period is 14 days.

In general there is no need to enter the greenhouse during the first 7 days after application of metam. However, if it is necessary, then respiratory protective equipment is required. Workers should also wear respiratory protective equipment when re-entering after 7 days to initiate ventilation of the treated structure (greenhouse or polytunnel). Re-entry to the greenhouse without respiratory protective equipment is possible after 14 days, provided the greenhouse is thoroughly ventilated prior to re-occupation.

### PERSONAL PROTECTION EQUIPMENT (PPE) IN CASE OF ENTRY INTO A TREATED FIELD DURING THE RE-ENTRY PERIOD

- Chemical resistant rubber gloves
- Chemical resistant rubber boots
- Cotton coverall
- Facial mask with active carbon filter type A (brown colour code) for gases and organic vapours with boiling point > 65°C

## 5.6. CLEANING OF MATERIAL AND DRUM DISPOSAL

After application, it may be necessary to dilute with water any product remaining in the tank – 1:100 – and apply it to the treated soil. Machinery should be cleaned of soil and debris before moving to another disinfection site or prior to storage.

In drip-irrigation application drip lines should be flushed with water after application but without over-saturating the treated soil.

Never reuse empty drums/containers for another application.

Treat empty drums/containers as hazardous waste.

**Drums:** rinse with clear water avoiding risk of ditch and water surface contamination and bring it to a certified collection centre.

**IBC** (Intermediate Bulk Containers of about 1000 L): recuperation assured by the container manufacturer.

#### PERSONAL PROTECTION EQUIPMENT (PPE) WHEN CLEANING MATERIAL AND DRUMS DISPOSAL

- Chemical-resistant rubber gloves
- Chemical-resistant rubber boots
- Hermetic coverall
- Eye or facial protection
- Facial mask with active carbon filter type A (brown colour code) for gases and organic vapours with boiling point > 65°C

## 5.7. REMOVAL OF PLASTIC FILM SEAL

### AND/OR SOIL AERATION

This is another critical phase where a small risk persists for residual MITC gas emission although it is expected that dissipation is almost complete after appropriate soil-sealing time.

This is of importance both with regard to potential discomfort for residents, especially when adverse wind direction is present at the time of aeration and for the workers that operate the seal removal.

Another option is the preliminary perforation of the film in different places to reduce the possibility of an overwhelming emission at the time of complete film removal (should the gases have not sufficiently dissipated).

It is also possible that plastic film tarp should stay in place as mulch in the case of a planting-through crop such as strawberry or lettuce. In that case holes need to be cut in the film at appropriate distances.

Forced soil aeration or pre-cultural tillage after soil disinfection by rotative tractor-driven device should not be deeper than the disinfected target layer in order to avoid mixing up of potential plant disease inoculum from untreated zone.

Whenever exposure of workers exists, appropriate PPE should be used.

#### PERSONAL PROTECTION EQUIPMENT (PPE) FOR REMOVAL OF PLASTIC FILM SEAL AND SOIL AERATION

- Rubber gloves and rubber boots
- Coverall
- Air purifying respirator is compulsory if sensory irritation is present\*

\* Sensory irritation trigger value: MITC concentration in the air > 0.6 ppm - see section 5.9.

Removed plastic film tarp should not be reused and must be discarded for destruction or recycling depending on the possibilities available locally.

## 5.8. CLEANING OF PERSONAL

### PROTECTION EQUIPMENT (PPE)

Follow manufacturers' instructions for cleaning/maintaining PPE.

If no instructions for washables exist, use detergent and hot water.

Keep and wash PPE separately from other laundry.


Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate and do not reuse them.

Do not transport contaminated clothing inside a closed vehicle unless stored in a sealed container.

## 5.9. GENERAL RULES ABOUT

### THE USE OF FILTER CARTRIDGES

Three main situations determine whether or not full-face masks should be worn with air purifying respirator (APR) filter cartridge(s) and their sensory irritation trigger values. At the third level operation should cease and handlers must leave. This is visualized in the following diagram:



Filter cartridges or canisters must be replaced:

- Whenever odour or sensory irritation from this product becomes apparent during use
- If the measured concentration of MITC is greater than 6 ppm
- At the end of each day's work period in the absence of any instructions or indications regarding service life

## 5.10. MONITORING RESIDUAL MITC

Monitoring for residual air or soil MITC concentration has two major advantages:

1. Checking the work/residents' environment for safety; 2. Checking for potential phytotoxicity risk after disinfection and soil aeration.

### 5.10.1. CHECKING THE WORK ENVIRONMENT FOR SAFETY

A distinction should be made between air concentration just above the soil or even the plastic film surface, at workers' inhalation height and in the surroundings for bystander/resident exposure. In the latter case measurements at different distances downwind from the treated plot should be made.

Two major measuring principles are available:

#### 5.10.1.1. PHOTO IONIZATION DETECTION (P.I.D.)

This sensitive quick-response electronic device is easy to use. By choosing the 10.6 eV uv- lamp version, the response factor for MITC is high compared to the possible response with other gaseous metam degradation compounds. The monitoring device should preferably be zeroed on top of the same but untreated soil.

Ambient air is pumped along the inside uv-lamp, molecules are ionized, collected on a flow-through electrode where they are discharged again. Changes in electrical current between electrodes are amplified and are a measure for gas concentration after calculation with the specific



response factor/correction factor. The lower this factor value the more sensitive the P.I.D. for the measured compound.

#### 5.10.1.2. DETECTION TUBES

Different manufacturers of safety and respiration protection equipment have developed specific tubes for MITC measurement.



Depending on manufacturers' instructions a volume of ambient air is pumped through a sealed glass gas reaction tube after inlet and exhaust (pump-side) tips are broken off. The gas reacts with a colouring agent sorbed on carrier material. Readings are made of the discoloured zone on a numeric MITC concentration scale.



The picture on the right shows a reading of about 20 ppm.



Each tube is for a single measurement and it takes some time to replace and perform the readings. Working with these disposables is of lesser interest when many measurements need to be performed.

### 5.10.2. CHECKING FOR POTENTIAL PHYTOTOXICITY RISK AFTER DISINFECTION AND SOIL AERATION

In this case both air monitoring and soil monitoring should be considered. Air monitoring could be important for adjacent crops. It is obvious that when checking conditions prior to soil disinfection the eventual presence of adjacent crops is known and measures are taken to avoid phytotoxicity.

For this purpose, a P.I.D. instrument is the most appropriate device that can be used. Soil gas monitoring after disinfection focuses on preservation of the new crop to be sown or planted.

After tarp removal and/or aeration the most classic test to perform is the so-called cress or lettuce seeds germination test.

The principle is simple: MITC-sensitive and quickly germinating plant seeds are exposed to soil samples originating from the disinfected plot. This can be in the gas phase or by direct sowing. Such tests have to some extent been standardized.

Some local distributors/contractors offer such phytotoxicity monitoring kits to their customers.

## PROCEDURE TO PERFORM A CRESS OR A LETTUCE SEED GERMINATION TEST

- 

1 With a trowel, dig into the treated soil to, or just below, the depth of application.
- 

2 Remove 2 to 4 small (25 - 50 g) treated soil samples, mix lightly, and immediately place a portion in an airtight jar so that fumes will not escape. Use glass jars with gas-tight lids.
- 

3 Prepare a similar jar with untreated soil (untreated check) for comparison.
- 

4 Sprinkle cress/lettuce seeds on the moistened surfaces of the soil and immediately re-close the jars.
- 


5 Keep the jars at 18 to 30°C; do not place in direct sunlight. Direct sunlight may kill the seed by overheating. Lettuce seed will not germinate in the dark.
- 

6 Inspect the jars for germination in 1 to 3 days. The soil is safe for planting if seeds in the treated jar germinate normally compared to the untreated check.


If phytotoxicity risk persists, it may be necessary to aerate the treated soil once more, wearing the appropriate PPE. A new test then has to be performed.

As an alternative, sowing or (trans)planting a few of the crops to be installed on the treated area can be made but this requires a longer period to observe an eventual residual phytotoxicity risk.

### PROCEDURE TO PERFORM A TOMATO TRANSPLANT TEST




1  
Transplant 5 to 10 succulent, fast-growing tomato seedlings into disinfected beds approximately 10 to 15 cm deep. If there is variation in the field, plant into the heaviest, wettest soil.



2  
Do the same in a non-disinfected area.



3  
Inspect the seedlings in 2 days for wilting or "root burn."



4  
If plants in the disinfected zone look the same as those in the nondisinfected zone, it is safe to plant.

# SUMMARY

## CHAPTER 5

### METAM SOIL DISINFECTION STEP-BY-STEP STEWARDSHIP

- Metam products, like many other plant protection products, need special attention and are subject to different safety rules at storage, handling and application level.
- Their corrosive and irritating properties and the generation of the gaseous MITC when applied to the soil, necessitate a range of different actions comprising:
  - Preliminary prospection of the disinfection site including assessing risks for bystanders and residents, checking of weather forecast and proper field preparation. This may result in postponing the application in the case of one or more adverse conditions.
  - Wearing of personal protection equipment appropriate to the type of operation being carried out.
  - Posting of no-entry signs prohibiting entry to the treated fields.
  - Respect of re-entry period.
  - Cleaning of material and safe disposal of empty drums.
  - Monitoring of residual MITC.



# ADDENDA

## ABBREVIATIONS

**APR:** air purifying respirator

**C x t:** concentration-time product

**F.I.F.:** fully impermeable film

**I.P.M.:** integrated pest management

**LDPE:** low density polyethylene

**MITC:** methyl isothiocyanate

**P.I.D.:** photo ionization detector

**PPE:** personal protective equipment

**SCBA:** self contained breathing apparatus

**SL:** miscible with water formulation

**T.I.F.:** totally impermeable film

**V.I.F.:** virtually impermeable film

**WHC:** water holding capacity

**Concentration x time product (C x t):** A numerical value expressed in  $\text{g} \times \text{hr}/\text{m}^3$  obtained by multiplying disinfectant concentration ( $\text{g}/\text{m}^3$ ) and the time (hrs) this concentration is maintained being a measure for biological activity.

**Disease:** Plant diseases are caused by fungi, bacteria and viruses.

**Dissipation:** Plant protection active compound dissipation after application is caused by chemical, physical or biological decomposition, degradation or metabolism. A soil applied compound is also dissipated by loss to the atmosphere (fumigants), leaching or by irreversible binding to soil particles.

**Dormancy:** Status of inactive seeds.

**Dosatron:** Type of dosing pump used in irrigation system allowing programmable supply of soil applied plant protection products and plant nutrients.

**Drip irrigation:** A water-saving technique to irrigate crops and to apply nutrients. The supply can be automated and installations can be used for soil disinfection.

**Fungicide (fungicidal):** Chemical compounds or biological organisms used to control plant parasitic fungi. Metam-based soil disinfectants have a broad fungicidal activity.

**Goose foot:** Horizontal blade in the shape of a goose foot mounted on soil labour machine lifting the soil with the possibility to carry liquid disinfectant injection pipes.

**Heap disinfection:** Disinfection can be used to disinfect soil heaps on impermeable floor (container, concrete floor or plastic film) in e.g. 25 cm thick layer and covered with gas-tight plastic film, for use in seeding containers or for potting soil.

## GLOSSARY

**Break through:** Timing where the process of gas or disinfectant diffusion starts through the plastic film tarpaulin or alternative soil sealing. This depends on film quality (composition; thickness), water condensation underneath the film, and temperature.

**Herbicide (herbicidal):** Chemical compounds used to control unwanted plants. Metam-based soil disinfectants are herbicidal for non-dormant weed seeds, seedlings and young weeds.

**Nematicide (nematicidal):** Chemical compounds or biological organisms used to control plant parasitic nematodes. Metam-based soil disinfectants are nematicides in contrast to granular cultural applied products which are nematostatics and which need repeated application.

**Personal protective equipment:** clothing, eye and respiratory protection

**Pest:** Plant pests are caused by insects, acari (mites), nematodes and slugs.

**Rotavator:** Rotating (with S- or L-shaped blades) soil labour device that crumbles and homogenizes soil (with soil applied products where appropriate).

**Seal (soil):** Technique to reduce disinfectant and soil humidity loss by compacting top soil layer with a roll, or by humidification of the top soil layer, or by installing a gas-tight plastic film tarpaulin, or by combination of two of them.

**Shank/Chisel:** Knife-shaped vertical blades mounted on soil labour machine cutting the soil with the possibility to carry liquid disinfectant injection pipes.

**SL-formulation:** miscible with water formulation.

**Soil disinfection:** Killing or reduction of soil-borne plant pathogenic or pest targets without completely sterilizing the soil.

**Solarization:** Technique used for soil disinfection in southern or tropical countries by covering soil with plastic film during several weeks to months in order to reach high soil temperatures also launching specific soil processes that can result to quite high degree of soil disinfection. Also the combination with chemical soil disinfection at reduced disinfectant rates has shown interesting results.

**Temperature inversion:** Temperature inversion is a meteorological situation where temperature increases in higher air layers with lower temperature in the lower layer creating a trap for pollutants by preventing their dilution into the atmosphere.

# LEGAL NOTICE

The information relating to the use of the product is given for information purposes only. No guarantee is provided to the client that the product is adapted to the specific use for which is intended by the client. The client should perform his own test to determine the suitability for a particular purpose. Information on the various application possibilities of our products is provided on the condition that the client should adapt the information to the specific conditions of their use and to the characteristics of other products with which they are mixed.

Metam CLR™ 42% and metam KLR™ 54% are "Restricted Use Pesticides" in the U.S. Regulations on metam differ between countries. It is therefore necessary to follow the local rules and to refer to the product label requirements as they may vary from the information provided herein. Local or national regulations prevail over the guidance provided in this manual.

46

Legal notice

---

## EASTMAN

### Eastman Corporate Headquarters

P.O. Box 431  
Kingsport, TN 37662-5280 U.S.A.

U.S.A. and Canada, 800-EASTMAN (800-327-8626)  
Other locations, +(1) 423-229-2000

[eastman.com/locations](http://eastman.com/locations)

Although the information and recommendations set forth herein are presented in good faith, Eastman Chemical Company ("Eastman") and its subsidiaries make no representations or warranties as to the completeness or accuracy thereof. You must make your own determination of its suitability and completeness for your own use, for the protection of the environment, and for the health and safety of your employees and purchasers of your products. Nothing contained herein is to be construed as a recommendation to use any product, process, equipment, or formulation in conflict with any patent, and we make no representations or warranties, express or implied, that the use thereof will not infringe any patent. NO REPRESENTATIONS OR WARRANTIES, EITHER EXPRESS OR IMPLIED, OF MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, OR OF ANY OTHER NATURE ARE MADE HEREUNDER WITH RESPECT TO INFORMATION OR THE PRODUCT TO WHICH INFORMATION REFERS AND NOTHING HEREIN WAIVES ANY OF THE SELLER'S CONDITIONS OF SALE.

Safety Data Sheets providing safety precautions that should be observed when handling and storing our products are available online or by request. You should obtain and review available material safety information before handling our products. If any materials mentioned are not our products, appropriate industrial hygiene and other safety precautions recommended by their manufacturers should be observed.

© 2024 Eastman. Eastman brands referenced herein are trademarks of Eastman or one of its subsidiaries or are being used under license. Non-Eastman brands referenced herein are trademarks of their respective owners.





[eastman.com/cropprotection](http://eastman.com/cropprotection)

Eastman Corporate Headquarters  
P.O. Box 431  
Kingsport, TN 37662-5280 U.S.A.

U.S.A. and Canada, 800-EASTMAN (800-327-8626)  
Other locations, +(1) 423-229-2000

[www.eastman.com/locations](http://www.eastman.com/locations)



**EASTMAN**

